



PICARBUTRAZOX GROUP U17 FUNGICIDE

PULL HERE TO OPEN

Vayantis®

syngenta®

Fungicide

A seed treatment product for protection against certain diseases of corn and soybean

<i>Active Ingredient:</i>	
Picarbutrazox*	36.0%
<hr/>	
<i>Other Ingredients:</i>	64.0%
<hr/>	
<i>Total:</i>	100.0%

*CAS No. 500207-04-5

Vayantis® is a flowable concentrate for seed treatment containing 3.3 pounds picarbutrazox per gallon.

1 quart Net Contents

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1635

EPA Est. 100-NE-001

Product of Japan

**SCP 1635A-L1 0321
4133142**



TABLE OF CONTENTS

- 1.0 FIRST AID**
- 2.0 PRECAUTIONARY STATEMENTS**
 - 2.1 Hazards to Humans and Domestic Animals**
 - 2.2 Personal Protective Equipment (PPE)**
 - 2.2.1 User Safety Requirements
 - 2.2.2 Engineering Controls
 - 2.2.3 User Safety Recommendations
 - 2.3 Environmental Hazards**

DIRECTIONS FOR USE

- 3.0 PRODUCT INFORMATION**
 - 3.1 Resistance Management**
- 4.0 APPLICATION DIRECTIONS**
 - 4.1 TANK MIXES**
- 5.0 ROTATIONAL CROP RESTRICTIONS**
- 6.0 RESTRICTIONS**
- 7.0 SEED CONTAINER LABEL REQUIREMENTS**
- 8.0 CROP USE DIRECTIONS**
 - 8.1 Corn
 - 8.2 Soybeans
- 9.0 STORAGE AND DISPOSAL**
- 10.0 CONDITIONS OF SALE AND LIMITATION OF LIABILITY**

1.0 FIRST AID

FIRST AID	
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<p style="text-align: center;">HOTLINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p>	

2.0 PRECAUTIONARY STATEMENTS

2.1 Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

2.2 Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton® ≥ 14 mils
- Shoes plus socks

2.2.1 USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

2.2.2 ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

2.2.3 USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

2.3 Environmental Hazards

Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities. Use is also permitted as an end-use seed treatment on agricultural establishments before planting as specified in the Specific Crop Use Directions. **DO NOT** use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be used in liquid or slurry treaters only.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

FAILURE TO FOLLOW THE DIRECTIONS FOR USE, PRECAUTIONS AND RESTRICTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notifications to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (WPS).

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves: barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, Viton ≥ 14 mils
- Shoes plus socks

3.0 PRODUCT INFORMATION

Vayantis is a seed treatment fungicide which provides early season protection against Pythium and Phytophthora causing seed rot and damping-off. Where rate ranges are shown, use the higher rate when disease pressure is expected to be severe.

3.1 Resistance Management

PICARBUTRAZOX	GROUP	U17	FUNGICIDE
---------------	-------	-----	-----------

For resistance management, Vayantis contains picarbutrazox, a Group U17 fungicide. Any fungal population may contain individuals naturally resistant to Vayantis and other Group U17 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies must be followed.

To delay fungicide resistance: Where possible, rotate the use of Vayantis or other Group U17 fungicides with different groups that control the same pathogens.

Vayantis is a U17 fungicide; therefore, continuous or sequential use of U17 fungicides within or during successive growing seasons must be avoided or reduced. Vayantis is not cross-resistant with other classes of fungicide that have different modes of action.

Seed treatment use must be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.

Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM information for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact Syngenta company representatives at 1-800-334-9481.

4.0 APPLICATION DIRECTIONS

Apply Vayantis as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of Vayantis into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed as stated in 40 CFR 153.155 (c). Allow seed to dry before bagging.

Follow the label directions for the most restrictive of label precautions and limitations. This product cannot be mixed with any product containing a label prohibition against such mixing.

4.1 TANK MIXES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Corn: For additional protection from certain seed-borne and soil-borne pathogens, Vayantis may be combined with seed treatment products containing mefenoxam, azoxystrobin, fludioxonil, sedaxane and/or thiabendazole. For insect protection, Vayantis may be tank mixed with seed treatment products containing thiamethoxam or cyantraniliprole. For nematode protection in addition to comprehensive fungicide defense and insect protection, Vayantis may be tank mixed with seed treatment products containing abamectin, thiamethoxam, thiabendazole, fludioxonil, mefenoxam, and azoxystrobin.

Soybeans: For additional protection from certain seed-borne and soil-borne pathogens, Vayantis may be combined with seed treatment products containing mefenoxam, azoxystrobin, fludioxonil, thiabendazole or sedaxane. For insect protection, Vayantis may be tank mixed with seed treatment products containing thiamethoxam or cyantraniliprole. For nematode protection in addition to comprehensive fungicide defense and insect protection, Vayantis may be tank mixed with seed treatment products containing pasteuria nishizawae or abamectin, and thiamethoxam, mefenoxam, fludioxonil, and/or sedaxane.

5.0 ROTATIONAL CROP RESTRICTIONS

In the event of crop failure or harvest of a crop grown from picarbutrazox treated seed, the field may be replanted immediately with labeled crops; all other crops - 30 days.

6.0 RESTRICTIONS

- **DO NOT** use for at-plant applications (e.g. hopper box, planter box, etc.). This product is to be used in liquid or slurry treaters only.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift.

7.0 SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with picarbutrazox fungicide.
- **DO NOT** use for feed, food, or oil purposes.
- User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seed treated with Vayantis:

- Store away from food and feedstuffs.
- **DO NOT** allow children, pets, or livestock to have access to treated seeds.
- Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. **DO NOT** contaminate water bodies when disposing of planting equipment washwater.
- Dispose of seed packaging in accordance with local requirements.
- In the event of crop failure or harvest of a crop grown from picarbutrazox treated seed, the field may be replanted immediately with labeled crops; all other crops - 30 days.
- Excess treated seed may be used for ethanol production only if:
 - (1) By-products are not used for livestock feed and
 - (2) No measurable residues of pesticide remain in ethanol by-products that are used for agronomic practice.
- For corn, **DO NOT** use at a rate that will result in more than 0.007 lb ai/A (3 grams ai/A) per year.
- For soybean, **DO NOT** use at a rate that will result in more than 0.017 lb ai/A (7.5 grams ai/A) per year.
- Vayantis treated seed may be planted on the same acres up to 2 times per year.

8.0 CROP USE DIRECTIONS

8.1 Corn

Crop	Diseases	USE RATE			
		fl oz/ 100 lb seed	fl oz/ 80,000 seeds unit	mg ai/seed	g ai/100 kg seed
Corn, includes field corn, sweet corn, popcorn and seed production	<i>Pythium</i> spp.	0.039 – 0.195	0.017 – 0.085	0.0025 – 0.0125	1 – 5
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.1. 					
The mg ai per seed, fl oz Vayantis per 100 lb seed, and fl oz Vayantis per 80,000 seeds rates are based on 1,800 seeds per pound.					
USE RESTRICTIONS					
1) DO NOT use at a rate that will result in more than 0.007 lb ai/A (3 grams ai/A) per year. 2) Vayantis treated seed may be planted on the same acres up to 2 times per year.					

8.2 Soybeans

Crop	Diseases	USE RATE			
		fl oz/ 100 lb seed	fl oz/ 140,000 seeds unit	mg ai/seed	g ai/100 kg seed
Soybeans	<i>Pythium</i> spp. <i>Phytophthora</i> spp.	0.039 - 0.195	0.018 – 0.090	0.0015 – 0.0075	1 – 5
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.1. 					
The mg ai per seed, fl oz Vayantis per 100 lb seed, and fl oz Vayantis per 140,000 seeds rates are based on 3,000 seeds per pound.					
USE RESTRICTIONS					
1) DO NOT use at a rate that will result in more than 0.017 lb ai/A (7.5 grams ai/A) per year. 2) Vayantis treated seed may be planted on the same acres up to 2 times per year.					

9.0 STORAGE AND DISPOSAL

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling (less than or equal to 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Container Handling (greater than 5 gallons)

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER

10.0 CONDITIONS OF SALE AND LIMITATION OF LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Vayantis®, the ALLIANCE FRAME,
the SYNGENTA Logo, and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



Viton® is a trademark of E. I. duPont de Nemours and Company, Inc.

©2021 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

**SCP 1635A-L1 0321
4133142**

For non-emergency (e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481



PICARBUTRAZOX GROUP U17 FUNGICIDE



syngenta

Fungicide

A seed treatment product for protection against certain diseases of corn and soybean



1 quart
Net Contents



(01) 0 07 02941 70893 8



Active Ingredient:	
Picarbtrazox*	36.0%
Other Ingredients:	64.0%
Total:	100.0%

*CAS No. 500207-04-5

Vayantis® is a flowable concentrate for seed treatment containing 3.3 pounds picarbtrazox per gallon.

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

EPA Reg. No. 100-1635

EPA Est. 100-NE-001

Vayantis®, the ALLIANCE FRAME, the SYNGENTA Logo, and the PURPOSE ICON are Trademarks of a Syngenta Group Company

©2021 Syngenta

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1635A-L1 0321
4133142

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOTLINE NUMBER: For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards: Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Handling (less than or equal to 5 gallons): Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER

