

Disease Risk Fungicide Schedule: East Texas



PEANUT FUNGICIDE PROGRAM

PLANTING DATE	At Planting	50–55 Days After Planting (DAP)	60 DAP	75 DAP	90 DAP	105 DAP	120 DAP
MODERATE RISK 5-6 Spray Program		Leaf Spot	Leaf Spot	S Blight / Limb Rot	Pythium	Pythium / S Blight / L Rot	Leaf Spot
PROGRAM Rate per acre	 2 fl oz per 100 lb/seed	 5.5 fl oz + 24.0 fl oz	 3.4 fl oz	 9.5 oz + 12.0 fl oz	 24.0 fl oz	 9.5 oz + 12.0 fl oz	 24.0 fl oz

Programs developed through the cooperation of



Key Diseases Controlled



	Trebuset® Peanuts	Alto 100SL	Bravo WeatherStik®	Miravis®	Quadris® / Abound®	Elatus®
Crown rot aspergillus	✓					
Cylindrocladium black rot	✓					S
Leaf spots	✓	✓	✓	✓	✓	✓
Pepper Spot			✓	✓	✓	✓
Pythium pod rot					S	
Rhizoctonia peg and pod rot					✓	
Rhizoctonia limb rot	✓	✓				✓
Rust		✓	✓		✓	✓
Seedling damping-off	✓					
Southern stem rot / White mold	S	✓		S	✓	S
Web blotch		✓	✓	✓	✓	✓

S = Suppression

© 2024 Syngenta. **Important: Always read and follow label instructions. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.** Abound®, Alto®, Elatus®, Miravis®, Quadris®, Trebuset® and the Syngenta logo are trademarks of a Syngenta Group Company. Bravo Weather Stik® is a trademark of ADAMA Corporation. Peanut Rx™ is a trademark of the University of Georgia. All other trademarks or service marks are the property of their respective owners. Use of third party trademarks are for informational use only and are not intended to imply endorsement by or affiliation with trademark owners.

GS 9095_3_1

SLC 15458A 09-2023



Scan the QR code to get more information about building the peanut program that's right for your field.

